


Sensitive and Terrestrial Ecosystems Labels



Sensitive Ecosystems Inventory of the Sunshine Coast and Adjacent Islands



- Legend for Sensitive and Terrestrial Ecosystems Labels, listing codes and descriptions for various ecosystem types.



Sensitive Ecosystems

Sensitive ecosystems are fragile and/or rare, or are ecologically important because of the diversity of species they support.

Old Forest (OF): Conifer-dominated dry to moist forest types, structural stage 7 (see table), generally <250yrs.

Woodland (WD): Dry open forests, generally between 10 and 30% tree cover, can be conifer dominated or mixed conifer and deciduous.

Herbaceous (HB): Non-forested ecosystems less than 10% tree cover, generally with shallow soils and often with bedrock outcroppings.

Riparian (RI): Areas adjacent to water bodies (rivers, lakes, ocean, wetlands) which are influenced by factors such as erosion, sedimentation, flooding and/or subterranean irrigation due to proximity to the water body.

Wetland (WN): Areas that are saturated or inundated with water for long enough periods of time to develop vegetation and biological activity adapted to wet environments.

Cliffs (CL): Very steep slope, often exposed bedrock, may include steep sided sand cliffs, habitat for rare species.

Other Important Ecosystems: Other important ecosystems have high biodiversity values.


Mature Forests (MF): Usually conifer-dominated, occasionally deciduous, dry to moist forest types, structural stage 6, generally >50yrs - >25 ha of buffering sensitive ecosystems.

Seasonally Flooded Agricultural Fields (FS): Annually flooded cultivated fields or hay fields; important migrating and wintering waterfowl habitat.

Other Mapped Ecosystems: Other mapped ecosystems occur in mosaic with sensitive ecosystems and are not possible to delineate separately at the mapping scale.

Young Forests (YF): Limited to areas of young forest dispersed among sensitive and other important ecosystems.


Polygon Label: Includes a field sample was completed but was not mapped.



Some polygon labels will have class and subclass repeated up to three times. This is not an error; it reflects the variability in site units and structural stages occurring within a polygon.

Ecosystem Components

This cartographic product uses Dot Density to indicate where more than one ecosystem class is mapped in a polygon.



Biogeoclimatic Units

- Geographic units: Coastal Douglas-fir Moist Maritime Subzone, Coastal Western Hemlock Eastern Very Dry Maritime Variant, Coastal Western Hemlock Dry Maritime Subzone, Coastal Western Hemlock Submontane Very Wet Maritime Variant.

Ecosystems

- Ecosystems: Georgia Lowlands Ecosystem, Strait of Georgia Ecosystem, Old Fieldland Ecosystem, Southern Pacific Ranges Ecosystem.

Map Symbols

- Map Symbols: Polygon Boundary, Biogeoclimatic Boundary, Ecosystem Boundary, Study Area Boundary, Roads, 20m contours, TRM Streams, Additional streams, Intermittent/Perennial Stream, Drainage Route.

Table with columns for Structural Stage, Substage, and description of various ecosystem types.

Table with columns for Terrestrial Ecosystem Map Codes and Site Unit Names, listing codes and their corresponding ecosystem names.

What is a Sensitive Ecosystem? For the purpose of this study, an ecosystem is considered to be a portion of the landscape with relatively uniform dominant vegetation.

Rationale: Ecologically significant lands and important wildlife habitats are fast disappearing throughout the lowlands surrounding the Strait of Georgia.

Purpose: The purpose of the Sensitive Ecosystems Inventory (SEI) of the Sunshine Coast is to identify, classify and map sensitive terrestrial ecosystems along the coastal lowlands.

Methodology: The mapping methods are based on the Vancouver Island SEI project and the Resources Information Standards Committee (RISC) Standard for Terrestrial Ecosystem Mapping (TEM) in BC.

Data Limitations: The SEI is a tool to alert decision makers to the existence of sensitive ecosystems, however when land-use changes are proposed detailed site-level assessments are necessary.

What can be done to protect sensitive ecosystems? Direct and indirect impacts to these ecosystems can be avoided by retaining or creating vegetated buffers around sensitive ecosystems to isolate them from outside disturbance.

Plan and implement all development activities in a manner that will not adversely affect or disturb the sensitive ecosystem.

A decision-maker (such as a politician or resource manager): ensure that protection of remaining sensitive ecosystems is a priority at all levels, and support programs, plans and operational activity that will help protect sensitive ecosystems.

A scientist: use your expertise to help identify sensitive ecosystems, define sites that need to be addressed, formulate conservation plans, contribute to the development of conservation and management strategies and explain to other professionals and decision makers the importance of sensitive ecosystems.

Acknowledgements: Environment Canada (Canadian Wildlife Service) and the B.C. Ministry of Sustainable Resource Management (MSRM) jointly managed this project.

Co-ordinator and extension: Carmen Cadrin, Justin Cullington, Jan Kirkby, Jo-Anne Stacey and Peggy Ward.

Field Crews: Louise Blight, Carmen Cadrin, Corey Erwin, Deepa South Pillay, Moraa Gray, Edwin Hubert, Stephen Hurlbut, Marc Johnson, Anne McIsaac, Wil Mackenzie, Claudia Schaefer, Jo-Anne Stacey and Leah Westergren.

