

---

# **Sensitive Ecosystems Inventory: Vernon Commonage 2005**

---

---

## **Volume 3: Wildlife Habitat Mapping**

---

---

**Allison Haney and Mike Sarell, Ophiuchus Consulting**

---



THE REAL ESTATE  
FOUNDATION  
OF BRITISH COLUMBIA



Canada



## Acknowledgements

---

The Steering Committee for the Vernon Commonage Sensitive Ecosystems Inventory has provided project management, guidance and direction throughout the project:

<b>Helen Davis</b>	<i>Project Coordinator</i>
<b>Carmen Wong</b>	<i>Allan Brooks Nature Centre</i>
<b>Susan Latimer</b>	<i>Ministry of Environment</i>
<b>Barbara Pryce</b>	<i>The Nature Conservancy</i>
<b>Susan Abbott</b>	<i>Greater Vernon Services Commission</i>
<b>Dale Danallanko</b>	<i>City of Vernon</i>
<b>Peter Tassie</b>	<i>North Okanagan Parks &amp; Natural Areas Trust</i>
<b>Christine LeFloch</b>	<i>Regional District of North Okanagan</i>
<b>Keith Louis</b>	<i>Okanagan Indian Band</i>
<b>Dale Donovan</b>	<i>Okanagan College</i>

The project was funded by The Real Estate Foundation of BC, the BC Ministry of Environment, the Habitat Stewardship Program<sup>1</sup>, the Greater Vernon Services Commission, and the Allan Brooks Nature Centre.

Project management and extension was provided by **Helen Davis**. Field work was completed by **Kristi Iverson**<sup>2</sup>, **Polly Uunila**<sup>3</sup>, and **Allison Haney**<sup>4</sup>. Bioterrain mapping was completed by **Polly Uunila** and ecosystem mapping was completed by **Kristi Iverson**. **Bon Lee**<sup>5</sup> completed digital and cartography work.

This project has adapted material from the reports for the Bella Vista – Goose Lake Range Sensitive Ecosystems Inventory and Central Okanagan Sensitive Ecosystems Inventory.

**Carmen Wong** provided accommodation for the field crew.

**Helen Davis** reviewed the draft version of this report.

We would like to thank the many landowners that gave us permission to access their lands for field sampling.

---

<sup>1</sup> The Habitat Stewardship Program for Species at Risk is a partnership-based, conservation initiative sponsored by the Government of Canada. The Program is managed cooperatively by Environment Canada, the Department of Fisheries and Oceans and Parks Canada, and is administered by Environment Canada.

<sup>2</sup> Iverson & MacKenzie Biological Consulting Ltd.

<sup>3</sup> Polar Geoscience

<sup>4</sup> Ophiuchus Consulting

<sup>5</sup> Baseline Geomatics Inc.

## Abstract

---

The Okanagan Valley contains the northern-most extent of Great Basin shrub-steppe ecosystems. These are often bisected by species-rich riparian and wetland habitats, and flanked by open forests and rugged slopes. The ensemble of wildlife that depends on habitats in the valley is diverse, containing species from the boreal forests to the north and the deserts to the south. Many of the southern-associated species are considered at risk in British Columbia and in Canada, due to their rarity and declining populations in landscapes that are sought for human development. In the North Okanagan, grasslands and shrub-steppe ecosystems dominate the lower elevations, and form the northern extent of these ecosystems in the valley. Extensive land development is fragmenting and encroaching on important wildlife habitats, contributing to wildlife and habitat declines.

This report is **Volume 3** of a Sensitive Ecosystems Inventory (SEI) project, initiated by the Allan Brooks Nature Centre. The report includes habitat summaries and species-habitat models for eleven wildlife species considered at risk in British Columbia. **Volume 1**<sup>6</sup> describes Sensitive Ecosystems, and offers practical advice on how to best avoid or minimize damage to them. **Volume 2**<sup>7</sup> provides details on the Terrestrial Ecosystem Mapping and terrain mapping.

The results of this habitat mapping indicate that abundant habitat exists for species dependant on rugged grasslands (e.g., Gopher Snake, Western Rattlesnake) and highly mobile species that require large expanses of open area (e.g., Swainson's Hawk, Badger). Habitat for species preferring certain grassland conditions such as gently sloping, large contiguous areas (e.g., Grasshopper Sparrow habitat) with low profile vegetation (e.g., Long-billed Curlew habitat) are scarcer. Although there is a fairly large amount of wetland habitat available for wildlife reliant on these habitats (e.g., Great Basin Spadefoot, Painted Turtle), there is a dearth of healthy riparian habitat, including mature to old deciduous forest habitat (e.g., Western Screech-owl habitat) and deciduous thickets with intact shrubby understory (e.g., Yellow-breasted Chat habitat). Overall, the mosaic of habitat types present in the study area leads to high habitat suitability for a wide range of wildlife species, and high biodiversity values.

Wildlife suitability models can be used alone to assess habitat values for individual species or in conjunction with Sensitive Ecosystems Inventory to identify potential environmental values of areas for conservation purposes (i.e., natural parks), or to guide development proposals. A Conservation Analysis has been conducted for this project, which should be used for landscape-level planning. For fine-scale evaluations, the wildlife models should be used to identify where environmental assessments should be conducted (areas with High and Moderate habitat suitability) if the lands are proposed for development. Environmental assessments for development proposals, including on-site inventory, should be conducted to verify and revise the predictive mapping. Revised environmental attributes in a georeferenced format can be returned to the planning staff at the City of Vernon or Regional District of the North Okanagan to revise in-house mapping. This would permit revisions to ecosystem and wildlife suitability mapping, updates of developed lands and areas retained as green space, and permit monitoring the efficacy of environmental planning and adaptive management.

---

<sup>6</sup> Iverson 2005

<sup>7</sup> Iverson and Uunila 2005

# Table of Contents

---

<b>ACKNOWLEDGEMENTS</b> .....	<b>III</b>
<b>ABSTRACT</b> .....	<b>IV</b>
<b>1 INTRODUCTION</b> .....	<b>1</b>
1.1 WHAT IS WILDLIFE HABITAT MAPPING? .....	1
1.2 HOW DOES WILDLIFE HABITAT MAPPING INTERACT WITH TEM AND SEI? .....	2
1.3 HOW IS WILDLIFE HABITAT MAPPING USED? .....	2
1.4 OBJECTIVES .....	3
<b>2 METHODS AND LIMITATIONS</b> .....	<b>3</b>
2.1 PROJECT WILDLIFE SPECIES .....	3
2.2 SPECIES-HABITAT MODELS.....	4
2.3 FIELD SAMPLING .....	5
2.4 WILDLIFE HABITAT MAPPING .....	5
2.5 MAPPING LIMITATIONS .....	6
<b>3 RESULTS</b> .....	<b>7</b>
3.1 SPECIES ACCOUNTS .....	7
3.2 FIELD SAMPLING .....	7
3.3 FINAL RATINGS TABLE .....	8
3.4 WILDLIFE HABITAT MAPS.....	9
<i>Great Basin Spadefoot</i> .....	10
<i>Painted Turtle</i> .....	11
<i>Western Rattlesnake</i> .....	12
<i>Gopher Snake</i> .....	13
<i>Swainson’s Hawk</i> .....	14
<i>Long-Billed Curlew</i> .....	15
<i>Western Screech-owl</i> .....	16
<i>Yellow-breasted Chat</i> .....	17
<i>Grasshopper Sparrow</i> .....	18
<i>Spotted Bat</i> .....	19
<i>Badger</i> .....	20
3.5 COMPOSITE CRITICAL HABITAT MAP.....	21
3.6 HABITAT VALUES OF SENSITIVE ECOSYSTEMS .....	22
<b>4 RECOMMENDATIONS</b> .....	<b>24</b>
4.1 GREAT BASIN SPADEFOOT .....	24
4.2 WESTERN RATTLESNAKE AND GOPHER SNAKE .....	25
4.3 LONG-BILLED CURLEW.....	25
4.4 SWAINSON’S HAWK .....	25
4.5 WESTERN SCREECH-OWL .....	25
4.6 GRASSHOPPER SPARROW .....	25
4.7 YELLOW-BREASTED CHAT .....	26
4.8 SPOTTED BAT .....	26
4.9 BADGER .....	26
<b>REFERENCES</b> .....	<b>27</b>
<b>APPENDICES</b> .....	<b>35</b>
APPENDIX A: DATA ACCESS .....	35
APPENDIX B: KNOWN AND POTENTIAL RARE WILDLIFE SPECIES IN THE STUDY AREA .....	36
APPENDIX C: WILDLIFE HABITAT ASSESSMENT FORMS .....	37
APPENDIX D: RATINGS TABLE.....	38

## List of Figures

---

Figure 1: Location of plots assessed during ecosystem mapping fieldwork.....	7
Figure 2: The many wetlands in the Commonage provide excellent breeding habitat for spadefoots.....	10
Figure 3: Distribution of suitable breeding and terrestrial habitats for Great Basin Spadefoot.....	10
Figure 4: Ponds provide living habitat for Painted Turtle.....	11
Figure 5: Distribution of suitable living and nesting habitats for Painted Turtle.....	11
Figure 6: Denning and basking habitat for rattlesnakes.....	12
Figure 7: Foraging habitat for rattlesnakes in the heat of summer.....	12
Figure 8: Distribution of suitable denning and foraging habitats for Western Rattlesnake.....	12
Figure 9: Warm aspect slopes with sparse tree cover and deep soils are important for egg laying and foraging for Gopher Snakes.....	13
Figure 10: Distribution of suitable denning, egg-laying, and living habitats for Gopher Snake.....	13
Figure 11: Expansive grassland for foraging and sporadic trees for nesting are critical for Swainson's Hawks.....	14
Figure 12: Distribution of suitable nesting and foraging habitats for Swainson's Hawk.....	14
Figure 13: Long-billed Curlews only nest on flat or gently sloping grasslands.....	15
Figure 14: Distribution of suitable nesting and rearing habitats for Long-billed Curlew.....	15
Figure 15: Birch and aspen provide the bulk of nesting opportunities, as cottonwood is scarce.....	16
Figure 16: Distribution of suitable nesting habitat for Western Screech-owl.....	16
Figure 17: A dense stand of rose and other deciduous shrubs provide potential nesting habitat.....	17
Figure 18: Distribution of suitable living (including nesting) habitat for Yellow-breasted Chat.....	17
Figure 19: Open grasslands with few shrubs are important characteristics of nesting habitats.....	18
Figure 20: Distribution of suitable living habitat for Grasshopper Sparrow.....	18
Figure 21: Crevices in large, sheer cliffs provide protection from predators.....	19
Figure 22: Distribution of suitable breeding habitat for Spotted Bat.....	19
Figure 23: Expansive, deep-soiled grasslands without road traffic are essential for Badger populations.....	20
Figure 24: Distribution of suitable living habitat for Badger.....	20
Figure 25: High and Moderate ratings for ten critical life requisites, displayed using highest value method..	21
Figure 26: Sensitive ecosystem mapping, displayed using largest area method.....	22
Figure 27: Conservation Zones resulting from the SEI Conservation Analysis.....	23

## List of Tables

---

Table 1: Habitat rating schemes for different knowledge levels of habitat requirements.....	2
Table 2: Wildlife species modelled in this project, their status, and rating scheme used.....	4
Table 3: Life requisites and habitat-uses rated during fieldwork.....	5
Table 4: Observations of project wildlife species or evidence of their use in the study area.....	8
Table 5: Map themes of habitat uses and life requisites modelled.....	9
Table 6: Map themes used in composite critical habitat map.....	21

# 1 Introduction

---

This report presents information on wildlife habitat mapping in the Vernon Commonage area between Okanagan Lake and Kalamalka Lake, including the Department of National Defence (DND) lands at the north end. It is the third volume in the Sensitive Ecosystems Inventory reports for Vernon Commonage.

**Volume 1**<sup>8</sup> describes inventory methods and results, rare and fragile ecosystems of the Vernon Commonage, highlights their values and importance, and offers practical advice on how to best avoid or minimize damage to them. **Volume 2**<sup>9</sup> provides details on the Terrestrial Ecosystem Mapping and terrain mapping.

## 1.1 What is Wildlife Habitat Mapping?

Habitat mapping portrays the potential importance of the land and its features to specific wildlife species through a species-habitat model. The model is used to generate a habitat map by assigning ratings to different habitat types, based on the needs of the species for particular life requisites. The ratings indicate the value of a habitat compared to the best habitat in the province<sup>10</sup>. Suitability is the ability of the habitat in its current condition to support a species. Capability is the ability of the habitat to support a species under optimal natural conditions, irrespective of the current condition of the habitat.

The following key elements and concepts summarize the Provincial standards for developing wildlife habitat ratings in British Columbia<sup>10</sup>:

1. There are three rating schemes; each reflects a different level of information available about the habitat requirements of a species (Table 1).
2. Ratings reflect a percentage of the provincial benchmark habitat. The provincial benchmark habitat has the highest suitability value for a given species in the province, against which all other habitats for that species must be rated. The benchmark is an actual location.
3. All ratings are a value for a specified season and activity, or life requisite.
4. A habitat rating is provided for each species for every occurring ecosystem unit (i.e., every site series / structural stage / site modifier combination).

Table 1 below shows the different habitat rating schemes.

---

<sup>8</sup> Iverson 2005

<sup>9</sup> Iverson and Uunila 2005

<sup>10</sup> Resources Inventory Committee 1999 (now Resources Information Standards Committee)

**Table 1: Habitat rating schemes for different knowledge levels of habitat requirements.**

Percent of Provincial Benchmark <sup>11</sup>	6-class (Substantial Knowledge of Habitat Use)		4-class (Intermediate Knowledge of Habitat Use)		2-class (Limited Knowledge of Habitat Use)	
76 - 100 %	High	1	High	H	Habitat Useable	U
51 - 75 %	Moderately High	2	Moderate	M		
26 - 50 %	Moderate	3				
6 - 25 %	Low	4	Low	L	Likely No Value	X
1 - 5 %	Very Low	5				
0%	Nil	6	Nil	N		

Habitat ratings are assigned to each ecosystem unit (e.g., habitat type) and then the values are projected onto the landscape where they are mapped. Habitat inventories assess the presence of available and potential habitat; they do not provide an indication of species presence or actual abundance. Much of the accuracy in predicting these habitat values is contingent on our understanding of how wildlife uses their habitats.

## 1.2 How does Wildlife Habitat Mapping interact with TEM and SEI?

Terrain and soil characteristics influence the vegetation of a site, within a given climate. Terrestrial Ecosystem Mapping (TEM) evaluates the specific ecological conditions (e.g. climate, terrain, vegetation community, and structural stage) for each polygon. All of these factors influence the wildlife assemblage and use within an area. TEM is used in a habitat model by assigning each ecosystem unit a wildlife habitat rating, indicating how useable (currently or potentially) the site is for a given wildlife species. These ratings are then applied to the TEM database and spatial data using GIS and portrayed as a habitat suitability or capability map of the study area.

In the field component of TEM the terrain, vegetation, and wildlife aspects are assessed in the field and discussed with the other members of the field crew, contributing to a greater accuracy of interpreted habitat use for wildlife. Field sampling is used to extrapolate the occurrence of certain habitat features as well, such as snags and coarse woody debris, to the types of habitats they commonly occur in.

Sensitive Ecosystems Inventory (SEI) takes into account ecological rarity and sensitivity of ecosystems, but also considers critical habitat needs for select wildlife species. Often, sensitive ecosystems contain important habitats for many wildlife species.

## 1.3 How is Wildlife Habitat Mapping Used?

The Okanagan Valley is one of the most diverse wildlife areas in Canada, and contains many of the Province and Nation's rare and endangered species. The area also has attracted considerable human settlement and associated land developments. Previous land development planning was limited in its ability to assess, identify, and conserve important wildlife habitats. This often led to the permanent loss of critical wildlife habitats, increasing the need to conserve those that remain. SEI and wildlife habitat

<sup>11</sup> The best habitat in the province. For example, High suitability (1 or H) is 76-100% as good as the best habitat in the province.

mapping can dramatically improve development planning to ensure that critical habitats are not developed, or that appropriate mitigation activities are undertaken.

The effectiveness of wildlife habitat mapping is contingent on the information being portrayed in a manner that is easily interpreted by planners, developers, regulatory agencies, and the public. This can be a challenge considering the diverse array of wildlife species potentially present, and the variety of habitat types used. Habitat values for wildlife have been considered to some degree in the SEI mapping, although 'Not Sensitive' ecosystems may still provide important habitat. Wildlife values for select species were given further consideration in the 'Conservation Analysis' provided in Volume 1<sup>12</sup>, which should be consulted for landscape-level planning. For land-use planning at a finer scale (e.g. neighbourhood plans), each species model should be inspected to direct detailed inventories to avoid or mitigate impacts to critical habitats.

Wildlife habitat mapping can also be used as a tool in wildlife management, a guide for wildlife viewing, and as a gauge of the loss of critical wildlife habitats.

## 1.4 Objectives

The objective of the wildlife habitat mapping is to provide input to land-use planning in the study area by providing estimated habitat values for wildlife species of management concern. The habitat mapping enables planners and managers to examine some of the wildlife values in order to guide development. Potential impacts can be identified and mitigation plans developed. ***Wildlife habitat mapping does not replace the need for development proponents to field-verify the presence of wildlife species and the significance of identified areas.***

## 2 Methods and Limitations

---

### 2.1 Project Wildlife Species

A vast number of rare or endangered wildlife potentially occur in the study area (Appendix B). Eleven of these wildlife species, all known to occur in the North Okanagan, were selected to demonstrate important wildlife habitats in the study area (Table 2). These species satisfy the following criteria<sup>13</sup> used to select wildlife species for habitat mapping:

- the level of knowledge of the species' use of habitat is adequate;
- the habitat required by selected species is also habitat required by other wildlife species;
- TEM is able to capture most of the habitat features required by the species;
- the species' habitat is present in the project area; and
- the species, or evidence of the species, is likely to be observed in the project area.

All of the selected species are considered at risk in the Province<sup>14</sup>, and some of these species have also been designated through Federal listing<sup>15</sup>. Species designated Threatened or Endangered are protected under the federal Species at Risk Act.

---

<sup>12</sup> Iverson 2005

<sup>13</sup> Resources Inventory Committee 1999 (now Resources Information Standards Committee)

<sup>14</sup> Conservation Data Centre (CDC) 2005: <http://srmwww.gov.bc.ca/cdc/>

<sup>15</sup> Committee on the Status of Wildlife in Canada (COSEWIC) 2005: <http://www.cosewic.gc.ca/>

**Table 2: Wildlife species modelled in this project, their status, and rating scheme used.**

Common Name	Scientific Name	Prov. Status <sup>16</sup>	COSEWIC Status <sup>17</sup>	Rating Scheme
Great Basin Spadefoot	<i>Spea intermontana</i>	Blue	Threatened	4-class
Painted Turtle	<i>Chrysemis picta</i>	Blue	-	4-class
Western Rattlesnake	<i>Crotalus oreganus</i>	Blue	Threatened	4-class
Gopher Snake	<i>Pituophis catenifer</i>	Blue	Threatened	4-class
Swainson's Hawk	<i>Buteo swainsoni</i>	Red	-	4-class
Long-billed Curlew	<i>Numenius americanus</i>	Blue	Special Concern	4-class
Western Screech-owl	<i>Megascops kennicottii macfarlanei</i>	Red	Endangered	4-class
Yellow-breasted Chat	<i>Icteria virens</i>	Red	Endangered	4-class
Grasshopper Sparrow	<i>Ammodramus savannarum</i>	Red	-	4-class
Spotted Bat	<i>Euderma maculatum</i>	Blue	Special Concern	4-class
Badger	<i>Taxidea taxus jeffersonii</i>	Red	Endangered	4-class

## 2.2 Species-Habitat Models

Wildlife habitat was modeled for the Vernon Commonage TEM according to the standards in the *BC Wildlife Habitat Ratings Standards - Version 2.0*<sup>18</sup>.

There are two basic components to a species-habitat model: the species account and the ratings table.

The species account summarizes the knowledge about a species and how it will be modeled. The account describes the distribution of the species in the province and in the project area, provides an overview of its ecology, and includes a detailed description of the critical life requisites and habitat uses of the species. The ratings section outlines the rating scheme (2, 4, or 6-class), the life requisites, and habitat uses that are modeled (map themes), and assumptions used to rate habitat characteristics. A section on map interpretation is also included, which describes how map themes were layered on the map, how the ratings were applied to the polygons, and provides information needed to correctly interpret each map.

Preliminary ratings tables, developed before field sampling, consist of an abbreviated table that provides habitat values for representative ecosystem units likely to occur in the project area. Our tables were modified to present assumptions used for rating ecosystems, which were incorporated into each species account. These assumptions, after being field-verified, guided development of the final ratings tables.

<sup>16</sup> Red List: indigenous species or subspecies (taxa) considered *Extirpated*, *Endangered*, or *Threatened* in BC.  
Blue List: indigenous taxa considered *Vulnerable* (Special Concern) in BC.

<sup>17</sup> Endangered = facing imminent extirpation in Canada or extinction.  
Threatened = likely to become endangered in Canada if limiting factors are not reversed.  
Special Concern = particularly sensitive to human activities or natural events.

<sup>18</sup> Resources Inventory Committee 1999 (now Resources Information Standards Committee)

## 2.3 Field Sampling

Field assessments occurred in conjunction with field sampling for ecosystem mapping. Survey intensity level 4 (visitation of 15 - 25% of polygons) was used<sup>19</sup>. Fieldwork took place in June of 2005. During field sampling, habitat values were recorded on Wildlife Habitat Assessment (WHA) forms (FS 882HRE 98/5). An example of the form is presented in Appendix C. Data was entered into Venus 5.0 data capture software. Table 3 lists and briefly describes the life requisites and habitat-uses rated in the field.

**Table 3: Life requisites and habitat-uses rated during fieldwork**

Species	Life Requisites and Habitat Uses	Rating Code
Great Basin Spadefoot	Security/thermal habitat for reproducing (breeding ponds).	RE
	Security/thermal habitat and food for general living, all year (terrestrial sites).	LIA
Painted Turtle	Security/thermal habitat for reproducing (egg-laying sites).	RE
	Security/thermal habitat and food for general living, all year (ponds).	LIA
Western Rattlesnake	Security/thermal habitat for general living all year (basking/denning sites).	LIA
	Food and security/thermal habitat for general living, summer.	LIS
Gopher Snake	Food and security/thermal habitat for general living, growing season.	LIG
	Security/thermal habitat for reproducing (egg-laying sites).	RE
Swainson's Hawk	Security habitat for reproducing.	RE
	Food for general living, growing season.	LIG
Long-billed Curlew	Security habitat for reproducing.	RE
	Food for general living, growing season.	LIG
Western Screech-owl	Security/thermal habitat for reproducing.	RE
Yellow-breasted Chat	Security/thermal habitat and food for general living, growing season.	LIG
Grasshopper Sparrow	Security/thermal habitat and food for general living, growing season.	LIG
Spotted Bat	Security/thermal habitat for reproducing (maternity roosts)	RB
Badger	Security/thermal habitat and food for general living, all year.	LIA

## 2.4 Wildlife Habitat Mapping

A final habitat ratings table was developed after field inspections were completed, and after a final list of ecosystem units was developed. Values were assigned using information from the species accounts, including assumptions, and from the wildlife report generated from field data in Venus 5.0.

We generated wildlife habitat maps by applying the ratings table values for each map theme (i.e., habitat use / life requisites for each species) to the TEM spatial and non-spatial data. An Ecosystem-based Resource Mapping (ERM) tool<sup>20</sup>, developed by the former Ministry of Sustainable Resource Management, was used to apply the ratings tables to the TEM map in ArcView GIS software.

<sup>19</sup> Resources Inventory Committee 1998 (now Resources Information Standards Committee)

<sup>20</sup> <http://srmwww.gov.bc.ca/wildlife/whr/sta.html>

Multiple map themes were displayed on the habitat-use map for some species, using a hierarchy of critical habitat requirements and life requisites. As habitat uses may overlap, we ensured that the most critical habitat uses overlaid less critical habitat uses. Each map was assigned a set of colours that identify the theme and values mapped.

Ratings were assigned to polygons with multiple ecosystem units (i.e., deciles) using one of the following four methods; based on which one best demonstrates the relative importance of that map theme:

- Highest-value – the highest rating within each polygon is displayed, regardless of the area it represents. The highest-value method exaggerates the amount of high value habitat because the whole polygon may be coloured high even if only a small part of it is actually high value.
- Averaged – the average rating within each polygon is displayed. Some parts of a polygon may be coloured as having some value, even if those parts have little or no habitat value. Similarly, some parts of a polygon may be rated as having low value, although the habitat in those parts has high value.
- Largest area – the rating for the ecosystem unit that covers the largest area of a polygon is displayed.
- Dot density – ratings for all of the ecosystems units are displayed, based on the percent area of the polygon they occupy. The dominant ecosystem unit provides the background colour, while dots of different colours or shades show the relative amount of other units occurring in the polygon.

## 2.5 Mapping Limitations

Limitations to Terrestrial Ecosystem Mapping are described in detail in Volume 1<sup>21</sup>, including:

- Scale of the aerial photographs (1:15,000). It is recommended that digital data not be enlarged beyond the scale of the photos as this may result in unacceptable distortion and faulty registration with other data sets.
- Date of the aerial photographs (1994) and field sampling (2005). On-going land uses may have changed some polygons after the date that the aerial photographs were taken or the field sampling was conducted.
- Ability to see disturbances such as cover of invasive plants on aerial photographs. Information from field sampling was applied to adjacent areas.
- Complex landscape, resulting in many complex polygons. Small ecosystems are often captured as a small component of a larger polygon that may contain up to three ecosystems.

For wildlife modelling purposes, additional limitations include:

- High variability of some ecosystem units (e.g., slope, soil depth, and, in a few units, vegetation composition). A given ecosystem unit may be described as having 'moderate to steep slopes', and some wildlife will use moderate slopes but are less likely to use steep slopes. Soil depth can also be highly variable; a shallow-soiled unit may have large pockets of deep soil suitable for burrowing.
- Condition of the habitat (e.g., understory fragmentation, forest ingrowth, invasive plants) is not accounted for in TEM, except for seral association in grasslands. This information is available in SEI as a condition value, and, while not incorporated into wildlife models, it was included in the Conservation Analysis<sup>22</sup>, where the sensitivity/rarity of the ecosystem, the condition of the ecosystem, and the wildlife values were all considered.

---

<sup>21</sup> Iverson 2005

<sup>22</sup> Volume 1: Iverson 2005

## 3 Results


---

### 3.1 Species Accounts

Complete species accounts, including citations, are available as described in Appendix A. Each species account also includes the final habitat suitability map for the species. Brief summaries of some important habitat requirements for the project species are included in the Wildlife Habitat Maps section below.

### 3.2 Field Sampling

A total of 169 plots were visited and assessed during Terrestrial Ecosystem Mapping and Sensitive Ecosystem Inventory, with 8 full plots, 69 ground inspections, and 92 visual inspections completed in the field (Figure 1). Only cursory investigations, if any, for evidence of wildlife use was conducted in some of the visual plots.



**Figure 1:** Location of plots assessed during ecosystem mapping fieldwork.

For many of the project wildlife species, we did not observe evidence of use during fieldwork. This is not surprising, as most of them are rare, elusive, or nocturnal, and fieldwork was intended as a habitat inventory rather than a wildlife survey.

Previous observation records for these species were amalgamated from various sources<sup>23</sup>. A summary of observations is presented in Table 4. Details of these observations are provided by species in Section 3.4. It should be noted that the relative abundance of species observations in the northern half of the study area is likely due, at least partly, to observer bias, with more records occurring in the area closer to Vernon and the Allan Brooks Nature Centre (ABNC).

**Table 4: Observations of project wildlife species or evidence of their use in the study area.**

Species	Previous Observations in Study Area	Observations During SEI
Great Basin Spadefoot	Many, MacKay Reservoir and north	None
Painted Turtle	Two locations, northeast portion	Many locations, throughout
Gopher Snake	Many, mostly on DND	None
Western Rattlesnake	Four known dens, east side	None
Swainson's Hawk	Several, all on DND	Two locations, just north of MacKay Reservoir
Western Screech-owl	None known (historically known at Okanagan Landing)	None
Long-billed Curlew	None	None
Grasshopper Sparrow	One location, DND	None
Yellow-breasted Chat	One location, Bailey Rd.	None
Spotted Bat	One location, above Kalamalka Lake	None
Badger	Several burrows, all on DND; one roadkill	One location, DND land

Other red- or blue-listed species recorded from the study area include Rubber Boa, Racer, California Gull, Lark Sparrow, and White-throated Swift.

### 3.3 Final Ratings Table

The final ratings table lists all of the mapped ecosystem units, including every combination of site series, site modifier, structural stage, stand modifier and seral association. See the expanded legend in Volume 3<sup>24</sup> for a description of all ecosystem units. Each ecosystem unit was assigned a rating for each of the 16 habitat uses for the eleven wildlife species. An example of the format of the ratings table is provided in Appendix D.

<sup>23</sup> CDC 2005, Ministry of Environment 2005, Clarke et al. 1993, Siddle 1993, Siddle 1995, Knopp et al. 2000, Sarell 2005

<sup>24</sup> Iverson and Uunila 2005.

### 3.4 Wildlife Habitat Maps

By applying the habitat ratings to the TEM database and spatial data, seventeen map themes were created (Table 5), including a duplication of one map theme (Gopher Snake denning uses the ratings from Western Rattlesnake denning).

**Table 5: Map themes of habitat uses and life requisites modelled.**

Species	Species Code	Map Themes	Rating Code
Great Basin Spadefoot	A-SPIN	Breeding General Living (foraging and burrowing)	RE LIA
Western Rattlesnake	R-CROR	Basking/denning Foraging	LIA LIS
Gopher Snake	R-PICA	Basking/denning <sup>25</sup> Foraging Reproducing (egg-laying)	LIW LIG RE
Swainson's Hawk	B-SWHA	Nesting Foraging	RE LIG
Long-billed Curlew	B-LBCU	Nesting Foraging	RE LIG
Western Screech-owl	B-WSOW	Nesting	RE
Yellow-breasted Chat	B-YBCH	General Living (nesting and foraging)	LIG
Brewer's Sparrow	B-BRSP	Nesting Foraging	RE LIG
Grasshopper Sparrow	B-GRSP	General Living (nesting and foraging)	LIG
Badger	M-TATA	General Living (denning and foraging)	LIA

The Species Accounts (see Appendix A) provide descriptions of how the map themes were rated and presented, as well as full-page maps for each species. Smaller versions of each map are presented in the following sections with an interpretation of each model. We discuss the distribution of habitats and the accuracy of the model based on past sightings and wildlife observations during fieldwork.

<sup>25</sup> Rattlesnake general living, all year (R-CROR\_LIA) ratings are used for this map theme.

## Great Basin Spadefoot

The Great Basin Spadefoot requires wetlands for courting, egg-laying, and development of eggs and larvae. The development of young spadefoots from egg to tadpole to adult is relatively quick, so temporary waterbodies that dry up in summer are commonly used. Ephemeral wetlands may actually be preferred due to the absence of fish or other aquatic predators.

Other than during spring breeding, adult spadefoots spend the rest of the year in nearby terrestrial habitats. These habitats must have deep, friable soils for burying themselves to avoid desiccation during dry weather and overwintering.

Many previous observation records exist for the northern portion of the study area, particularly DND land. They have also been recorded from MacKay Reservoir and northeast of Rose's Pond. Spadefoots were not detected during fieldwork; however, high suitability breeding ponds (Figure 2) appear to occur throughout the Commonage.




**Figure 2:** The many wetlands in the Commonage provide excellent breeding habitat for spadefoots.

The suitability model generated two map themes: aquatic breeding habitats and terrestrial living habitats (Figure 3). Breeding habitats overlay living habitats. Both themes were displayed using the highest-value method.

Suitable breeding sites predicted by the model occur throughout the study area, but they may not be used if suitable terrestrial habitats are not present (e.g., Predator Ridge). However, low rated terrestrial habitats near breeding ponds may be used to a higher extent than the rated value indicates, due to their proximity to breeding habitat. Conversely, high suitability terrestrial habitats may not be used if they are situated too far from breeding habitats, but it should be noted that very small, temporary 'wetlands' may not have been identified in the TEM.

Spadefoots are well adapted to desert conditions, with a hardened 'spade' on their hind foot for burrowing into soils, and skin secretion that prevents dehydration while buried.



**Figure 3:** Distribution of suitable breeding and terrestrial habitats for Great Basin Spadefoot.

## Painted Turtle

Turtles require wetlands throughout the year for foraging and over-wintering. Females leave the ponds to lay eggs in nearby terrestrial habitats with coarse, well-drained soils and sparse vegetation.

Turtles only leave their ponds when females lay eggs during the summer, and the occasional dispersal, particularly if their pond dries up during a dry spell.

Previous observation records exist for two locations in the northern portion of the study area: near the south end of DND land, and another pond further south.

Painted Turtles were detected at about a dozen ponds during fieldwork, and high suitability ponds (Figure 4) occur throughout the Commonage.




**Figure 4:** Ponds provide living habitat for Painted Turtle.

The suitability model generates two map themes: aquatic living habitats and terrestrial nesting habitats (Figure 5). Both themes are displayed using the highest-value method. Only nesting habitats within 150 m of suitable ponds are portrayed.

Suitable habitat predicted by the model occurs throughout the study area, particularly the central portion. Although the Commonage appears to support abundant turtle populations, high mortality may be occurring from road traffic and disturbance of nesting sites in exposed sand or gravel.


Turtles spend the winter in the mud at the bottom of ponds. During this period of inactivity, turtles respire by absorbing oxygen from water they take into their pharynx and cloaca (i.e., both ends of the digestive tract).



**Figure 5:** Distribution of suitable living and nesting habitats for Painted Turtle.

## Western Rattlesnake

Western Rattlesnakes require sparsely vegetated ecosystems such as rock outcroppings for hibernating. Riparian areas, broadleaf woodlands, grasslands, or open forests are used for foraging. High-value denning and basking habitats on south-facing rocky hillsides (Figure 6) were observed at eight of the field plots.



**Figure 6:** Denning and basking habitat for rattlesnakes.

High-value foraging habitats include riparian areas and broadleaf woodlands, which support dense prey populations and have more moderate summer temperatures (Figure 7).




**Figure 7:** Foraging habitat for rattlesnakes in the heat of summer.

Suitability was modeled for two map themes for rattlesnakes; both were displayed by the highest-value method (Figure 8). The denning theme (top map layer) consists of security/thermal habitats potentially used all year, including denning during winter, basking in spring and fall, and throughout the summer for gravid females. Foraging includes habitats that likely provide security and thermal shelter as well as food.

The map depicts suitable habitat throughout the study area, although they have only been recorded from the east side.

Much of the predicted denning in the southwest portion may be too treed (shaded) to provide the warmth required for winter hibernacula.

Rattlesnakes are the only dangerously venomous snake species in BC, but will rarely bite unless threatened.



**Figure 8:** Distribution of suitable denning and foraging habitats for Western Rattlesnake.

## Gopher Snake

Gopher Snakes den in either deep-soiled grasslands or sparsely vegetated ecosystems (rocky habitats). Deep soiled denning sites were not modeled for this project, as they are very difficult to predict. Because of the similarities in rocky den sites to rattlesnake suitability, ratings were not assigned separately for Gopher Snake.

High value foraging habitat occurs in deep-soiled grasslands, broadleaf woodlands and riparian areas.

Unlike Western Rattlesnakes, Gopher Snakes lay eggs. Egg-laying habitat is frequently associated with warm-aspect grasslands with deep soils (Figure 9). We assessed seven plots of the 169 with high-value egg-laying habitat.

Gopher Snakes have been previously recorded from numerous locations on DND land, and one location along Commonage Road north of MacKay Reservoir.




**Figure 9:** Warm aspect slopes with sparse tree cover and deep soils are important for egg laying and foraging for Gopher Snakes.

The Gopher Snake habitat-suitability model generated three map themes. Denning is the top map layer and overlays egg-laying, which overlays general living (Figure 10). Denning was derived from the rattlesnake denning theme, and predicts only rocky den sites. This model does not attempt to predict earthen burrows that may also be used by Gopher Snakes for over-wintering. Deep-soiled, warm aspect sites were used to predict egg-laying habitat, which may also capture some denning sites. The living theme depicts areas potentially rich in prey that also provide security and thermal cover.

Gopher Snakes likely occur throughout the study area, although the south-western portion may be too heavily forested to support large populations.

Although they resemble the rattlesnake, Gopher Snakes are constrictors, and non-venomous.



**Figure 10:** Distribution of suitable denning, egg-laying, and living habitats for Gopher Snake.

## Swainson's Hawk

These hawks require expansive, open areas for foraging, and scattered large trees in or adjacent to grasslands for nesting (Figure 11).

Swainson's Hawks are known from the northern part of the study area, and were observed foraging and roosting near MacKay Reservoir during fieldwork. They were seen at the north end of the District of Lake Country during the summer of 2005 as well.

Fifteen of 169 plots were assessed as having high value nesting habitat, and 27 as high-suitability for foraging, which indicates that abundant habitat exists.




**Figure 11:** Expansive grassland for foraging and sporadic trees for nesting are critical for Swainson's Hawks.

Both the nesting (top layer) and foraging theme generated by the model were displayed using the highest-value method (Figure 12).

Most of the nesting habitat depicted occurs in the southern portion of the study area. However, the small stands of trees and isolated trees within grassland habitats typical of the northern portion are likely more suitable for nesting.

Hawks are highly motile, hunting over a large area, and require a relatively large amount of suitable foraging habitat to support a nesting pair. Because of the availability of habitat, the study area likely has one of the highest concentrations of Swainson's Hawks in the province.

The colouration of Swainson's Hawks, as well as the more common Red-tailed Hawk, is highly variable. They can be distinguished by their longer, narrower, and more pointed wings.



**Figure 12:** Distribution of suitable nesting and foraging habitats for Swainson's Hawk.

## Long-Billed Curlew

Curlews require fairly large areas of level to gently sloping grassland with short vegetation and no trees for nesting. Families of curlews will often move to lush cultivated fields once the young have fledged. Foraging occurs in hayfields, pastures, meadows, and grasslands.

No sign of Long-billed Curlews was detected during fieldwork, and they have not been previously recorded from the study area.

High suitability nesting habitat (Figure 13) was encountered at only three plots during fieldwork. Expanses of gently sloping grasslands are typically the first areas to succumb to development pressures.




**Figure 13:** Long-billed Curlews only nest on flat or gently sloping grasslands.

The suitability model for curlews generates two map themes: nesting and foraging (Figure 14). Curlews generally avoid nesting near treed areas, so only polygons that contain 20% or less forested ecosystems are displayed.

A fair amount of predicted high suitability habitat appears on the map. Much of this area, however, may not fulfill curlews' preference for large, open and flat areas with low-profile vegetation. Despite the availability of grasslands in the study area, optimum nesting conditions are scarce due to slope or proximity to trees. The central portion of the study area and the DND lands appear to have the highest potential.

Curlews are very tolerant of cattle grazing, except that they are vulnerable to trampling of the eggs and young.



**Figure 14:** Distribution of suitable nesting and rearing habitats for Long-billed Curlew.

## Western Screech-owl

Western Screech-owls are dependant on mature to old riparian forest and most often nest in cavities in large cottonwood trees. Nesting is known from the Okanagan valley floor as far north as Coldstream Creek, and also in the middle Shuswap (J. Hobbs, H. Davis pers. comms.).

We found no evidence of Western Screech-owls during fieldwork, and no previous records exist for the study area. Historical records do occur at Okanagan Landing.

Potential high-value nesting habitat was observed at only two plots (COMG007 and 020669), both were dominated by large paper birch. A number of aspen gullies were assessed as moderate suitability (six plots; Figure 15).




**Figure 15:** Aspen and birch provide the bulk of nesting opportunities, as cottonwood is scarce.

The suitability model for Western Screech-owl generates only one map theme, nesting habitat, which is displayed using the highest-value method (Figure 16). Some foraging may occur in adjacent areas.

There is very little suitable habitat predicted for screech-owls, and inventories are required to determine whether they are present in the study area. The riparian areas on the Commonage tend to be small in nature, and many have been impacted by land practices. Optimal habitat for screech-owls (mature cottonwood stands) more commonly occurs in the valley bottom.

Most of the portrayed suitable habitats are mature forest or riparian ecosystems, with mixed coniferous and deciduous overstories.

The call of the Western Screech-owl is easily identified, described as a 'bouncing ping-pong ball'.



**Figure 16:** Distribution of suitable nesting habitat for Western Screech-owl.

## Yellow-breasted Chat

These songbirds are dependant on riparian areas with a shrubby understory, preferably with dense wild rose and snowberry.

Yellow-breasted Chats were not observed during fieldwork. One record exists from the study area (on Bailey Road).

High suitability habitat for Yellow-breasted Chats (Figure 17) was recorded at five plots. Many other sites would be of high value except that the amount of cattle use has resulted in degradation of the shrubby understory vegetation.




**Figure 17:** A dense stand of rose and other deciduous shrubs provide potential nesting habitat.

All chat activity is generally confined to a nesting territory. Therefore, there is only one map theme (living), which includes nesting and foraging (Figure 18). This theme is displayed using the highest-value method.

Chat habitat often occurs as small strips or pockets, and likely occupies only a portion of some of the polygons identified. These are usually located in gullies or around wetlands.

Chats earned their name because of their noisy and highly diverse range of calls, including a typical 'chat-chat-chat-chat'. They are one of the very few songbirds that are vocal at night.



**Figure 18:** Distribution of suitable living (including nesting) habitat for Yellow-breasted Chat.

## Grasshopper Sparrow

Grasshopper Sparrows generally occur in grasslands with little or no sagebrush, which are flat or on gentle warm aspects.

Although not detected during fieldwork, Grasshopper Sparrows have been recorded from one location on DND lands, and are known to breed regularly around Goose Lake west of Vernon.

High suitability nesting/foraging habitat (Figure 19) was encountered at a large number of the plots assessed.




**Figure 19:** Open grasslands with few shrubs are important characteristics of nesting habitats.

Nesting and foraging by Grasshopper Sparrows generally occurs in the same type of habitat. Therefore, the model generated only one map theme: living (Figure 20). The theme is displayed using the dot-density method, as this bird prefers fairly large areas of suitable habitat. This allows the visualization of contiguity and where unsuitable habitats occur in otherwise suitable polygons.

Large areas of high-rated living habitats were scarce but concentrated in the north end of the study area. High and moderate rated living habitats should be the target of inventories.

Grasshopper Sparrows nest on the ground, usually at the base of bunchgrasses, and use the overhanging vegetation to build a dome with a side entrance. They received their name from a portion of their call that resembles the buzz of a grasshopper.



**Figure 20:** Distribution of suitable living habitat for Grasshopper Sparrow.

## Spotted Bat

Spotted Bats roost in crevices in large, sheer cliffs, which are also used by maternal colonies where females give birth to young.

Only one roost location is known from the study area, in the cliffs above Kalamalka Lake across from Cosens Bay.

No high suitability habitat was encountered during fieldwork (the known roost site was not sampled). Only one moderate suitability cliff was observed, at the south end of DND land (Figure 21).




**Figure 21:** Crevices in large, sheer cliffs provide protection from predators.

The Spotted Bat suitability model generates just the one theme: breeding, which also includes non-maternity roosting (Figure 22).

The model predicts very little suitable habitat, as verified by fieldwork. Because of their scarcity, the suitable cliffs that do exist are extremely important for this species.

Spotted Bats are the only bat species in BC whose echolocation calls are audible to the human ear, which sound like a series of high-pitched ticks.



**Figure 22:** Distribution of suitable breeding habitat for Spotted Bat.

## Badger

Badgers are usually residents of deep-soiled grasslands (Figure 23) although they will venture into a broad range of habitats. The north Okanagan has an abundance of deep-soiled grasslands that probably historically supported stable Badger populations.

We found old Badger burrows at one location in a man-made ridge on DND land, where digging would be easy because of the lack of soil compaction.

Many plots were assessed as high-value habitat during fieldwork, including suitability for maternity dens.




**Figure 23:** Expansive, deep-soiled grasslands without road traffic are essential for Badger populations.

One map theme, living, is generated by the model, which includes foraging and denning (Figure 24). The dot density method is used to display habitat values, as this gives an indication of the proportion of the polygon suitable for use.

The abundance of rodent prey could not be directly included in the habitat suitability model. Pocket gopher burrows often occurred in small pockets of deep soil throughout the rolling topography of much of the study area. However, badgers commonly forage for more colonial prey (i.e., marmots and ground squirrels), displaying patchy use of habitats.

Badger populations have likely declined from habitat loss, persecution and traffic mortality. Fragmentation of habitats has also likely contributed to their decline. The study area and the Bella Vista – Goose Lake Range are important refuges of expansive grasslands suitable for Badgers.



**Figure 24:** Distribution of suitable living habitat for Badger.


### 3.5 Composite Critical Habitat Map

Ten life requisites were chosen to represent the most limiting habitat requirements of the project wildlife species (Table 6). This does not imply that the species or life requisites omitted are not as important. Rather, their needs may be met if habitat for the remainder of the map themes is conserved.

**Table 6: Map themes used in composite critical habitat map.**

Species	Species Code	Map Themes	Rating Code
Great Basin Spadefoot	A-SPIN	Breeding	RE
Western Rattlesnake	R-CROR	Basking / denning	LIA
Gopher Snake	R-PICA	Egg-laying	RE
Swainson's Hawk	B-SWHA	Nesting	RE
Long-billed Curlew	B-LBCU	Nesting	RE
Western Screech-owl	B-WSOW	Nesting	RE
Yellow-breasted Chat	B-YBCH	General Living (nesting and foraging)	LIG
Grasshopper Sparrow	B-GRSP	General Living (nesting and foraging)	LIG
Spotted Bat	M-EUMA	Breeding/roosting	RB
Badger	M-TATA	General Living (denning and foraging)	LIA

A composite critical habitat map of high- and moderate-value habitats for the ten critical map themes was generated and is presented in Figure 25. This map is displayed using the highest-value method. While this method is excellent for highlighting polygons containing important areas, it often tends to exaggerate the amount of valuable area, as entire polygons are shown by the highest value they contain.




**Figure 25:** High and Moderate ratings for ten critical life requisites, displayed using highest value method.

The composite wildlife map portrays abundant high-suitability habitat, indicating that the majority of polygons in the study area contains valuable habitat for at least one of the project species. The map should be used to view important habitats on a landscape level. For areas of interest, refer to individual wildlife habitat models and investigate them in the field to assess values.

### 3.6 Habitat Values of Sensitive Ecosystems

Sensitive Ecosystem Inventory categories<sup>26</sup> are shown in Figure 26 by largest area, which portrays the dominant component of each polygon. Almost all polygons dominated by *sensitive ecosystems* have high suitability for at least one of the project wildlife species (see Figure 25). *Other important ecosystems*, particularly disturbed grasslands, often have high value for many of the project wildlife species as well. It should be noted that because the SEI categories are displayed using largest area, many of the polygons likely contain higher SEI values than shown.




Many polygons without sensitive or other important ecosystems may still provide important wildlife habitat for species at risk, including rural and agricultural areas, and very weedy grasslands with little or no native vegetation.

**Figure 26:** Sensitive ecosystem mapping, displayed using largest area method.

<sup>26</sup> Iverson 2005

The Conservation Analysis described in Volume 1<sup>27</sup> takes into account not only the rarity and fragility of ecosystems (sensitive ecosystems), but also the condition of the ecosystems and wildlife values (Figure 27). The Conservation Zones resulting from the Conservation Analysis appear to protect the bulk of critical habitat for all project species, including important wildlife corridors.



**Figure 27:** Conservation Zones resulting from the SEI Conservation Analysis.

---

<sup>27</sup> Iverson 2005

## 4 Recommendations

---

Local government, BC Parks, landowners, consultants, and other interested groups can use the wildlife habitat mapping in a number of ways. As a management tool, the wildlife suitability maps can be used to direct broad wildlife management strategies, such as recovery of habitats for species at risk and ecosystem management practices, including prescribed burns. As a landscape-level planning tool, the Conservation Zones (Figure 27) resulting from the Conservation Analysis can be used to direct development towards less sensitive areas. The composite critical habitats map (Figure 25) should be used to identify potentially critical areas that should be considered for conservation unless an environmental impact assessment recommends adjustments to these boundaries. A development permit bylaw could restrict development on these areas until they are assessed. Assessments should address the relevancy of each of the wildlife suitability models within the area of interest, as a minimum standard. A useful template of Terms of Reference can be found in the Habitat Atlas for Species at Risk<sup>28</sup>. Volume 1<sup>29</sup> of the Sensitive Ecosystem Inventory contains additional environmental impact assessment guidelines.

Due to the wildlife significance of the area, environmental impact assessments should not only concentrate on ground-truthing the results of these suitability models, but should also inventory for other species at risk and their critical habitats. Volume 1<sup>29</sup> provides lists of species at risk that may be associated with each sensitive or other important ecosystem.

Anyone conducting environmental impact assessments using this information should have a good understanding of each species' habitat requirements and associated threats when evaluating development impacts and establishing environmentally sensitive areas (ESA). Best Management Practices are being developed for many species at risk, and these should be consulted in addition to the management recommendations outlined here.

Many wildlife species require connectivity throughout their range, and this should be given consideration when assessing the lands of interest in context with the surrounding area. Priority areas should be covenanted or otherwise designated for conservation.

The following are brief management guidelines for each of the project wildlife species.

### 4.1 Great Basin Spadefoot

Inventories are required to determine which ponds are used for breeding. This data can be used to adjust the suitability for terrestrial habitats. Generally, buffers around breeding sites should be at least 350 m<sup>30</sup> to protect both breeding and adjacent terrestrial habitats and to avoid road and other mortality. However, this could vary depending on the suitability of upland habitat. Spadefoots may travel several hundred metres from ponds, and up to 1.5 km, so buffers should be extended to encompass the highest-suitability surrounding habitat, attempting to capture at least 5 ha of terrestrial area<sup>31</sup>.

Corridors must be maintained between ponds and foraging sites. Developments that pose a hazard or obstruction to spadefoots, including roads, retaining walls, and steep-sided trenches, should not occur between aquatic breeding habitats and nearby suitable terrestrial habitats. Management should also

---

<sup>28</sup> BC Environment 1998, pg 108

<sup>29</sup> Iverson 2005

<sup>30</sup> Semlitsch and Bodie 2003

<sup>31</sup> Sarell 2004

consider the connectivity between aquatic habitats, to maintain gene flow between spadefoot populations. Artificial breeding habitats can be created as part of mitigation programs.

## **4.2 Western Rattlesnake and Gopher Snake**

Management of Low, Moderate and High potential denning habitats should include a no-development zone, unless an inventory has demonstrated that the depicted habitat(s) are not used. Recreational corridors should avoid these areas to minimize human-snake conflicts, including mortality from mountain bikes and vehicles. Summer foraging areas should be carefully assessed to determine whether any development is appropriate, and if so, what mitigation measures are required. Although corridors to allow snake movement from winter security/thermal habitats to summer foraging habitats have not been mapped, they should be interpreted and applied to project planning. Roads should not intersect any of these areas unless appropriate mitigation measures are employed to avoid traffic mortalities. Paved roads are a particularly large threat to snakes due to their habit of basking on the warm surface for thermoregulation. Snake exclusion fencing may be required to reduce encounters and mortality in developed areas.

## **4.3 Long-Billed Curlew**

Conduct inventories in grassland habitats during the breeding season to determine whether Long-billed Curlews are present. Curlews require an expanse of level to gently sloping grasslands. Any development in these areas, including roadways and recreational corridors, will significantly impact these birds. Livestock should not access these areas during the breeding season to protect nests from trampling. Domestic cats should not be permitted in these areas as they may prey upon adults and nestlings.

## **4.4 Swainson's Hawk**

Inventories during the breeding season should be conducted to locate existing nest trees. Conserve wide grassland networks between nest trees and other suitable nesting habitats. Do not locate transportation or recreational corridors within 100 m of nest trees.

## **4.5 Western Screech-owl**

Spring inventories are required to determine whether nesting occurs in riparian forests in the study area. Maintain deciduous and mixed stands, including wildlife trees, to provide nesting and foraging habitats. Incorporate surrounding natural habitats, particularly meadows, as a buffer to these areas. Nest boxes can help to mitigate small losses of nesting habitat.

## **4.6 Grasshopper Sparrow**

Breeding season inventories are required to determine the extent to which they occur in grassland habitats, including weedy sites. They are semi-colonial but often shift their breeding territories between years. Therefore, additional suitable grassland habitats should be retained to accommodate breeding in subsequent years. A buffer to reduce disturbances is also recommended. Livestock should not access these areas during the breeding season to protect nests from trampling. Domestic cats should not be permitted in these areas as they may prey upon adults and nestlings.

#### **4.7 Yellow-breasted Chat**

Inventories during the breeding season are required to determine where they occur in the study area. Maintain deciduous stands and restore shrubby understory, particularly wild rose. Livestock should have limited access to these areas as they reduce the shrubby component of these ecosystems. Buffers should be incorporated to reduce disturbances to these areas. Domestic cats should not be permitted in these areas, as they may prey upon adults and nestlings.

#### **4.8 Spotted Bat**

Spotted Bats roost in large cliffs and may hibernate in these features as well. Generally there are few impacts to cliffs from human activities. Development and blasting should not be permitted within 200 m of a roost cliff. New developments should have shielded streetlights. Recreational rock climbing should not be permitted on roost cliffs.

#### **4.9 Badger**

Inventories should be conducted to locate burrows, particularly maternal burrows, although differentiating between maternal and other types of burrows is difficult. The most critical habitat sites for Badgers are their maternal dens and adjacent foraging areas. Burrows usually occur in deep soils on gentle to moderate sloping grasslands, often adjacent to significant populations of ground squirrels, marmots or pocket gophers. Management should ensure there is no disturbance to occupied or maternal burrow sites and that no activities significantly affect prey species or create barriers between suitable areas. Corridors or connectivity should be maintained with other natural areas to allow for their high degree of motility and dispersion. Road placement should avoid intersecting suitable badger habitat, as road mortality is the major cause of death for this species (Weir et al. 2005). Landowners may wish to conduct inventories to specifically identify important badger habitats.

## References

---

- Adams, I., T. Antifeau, M. Badgry, L. Campbell, A. Dibb, O. Dyer, W. Erickson, C. Hoodicoff, L. Ingham, A. Jackson, K. Larsen, T. Munson, N. Newhouse, B. Persello, J. Steciw, J. Surgenor, K. Sutherland and R. Weir. 2003. National recovery strategy for the American badger, *jeffersonii* subspecies (*Taxidea taxus jeffersonii*). Recovery of Nationally Endangered Wildlife (RENEW), Ottawa, Ontario.
- Adams, I.T. and T.A. Kinley. 2003. Badger (*Taxidea taxus jeffersonii*), in Identified Wildlife management strategy - accounts and measures for managing Identified Wildlife, Version 2004. Ministry of Water, Land and Air Protection, Victoria, BC.
- Allen, J.N. 1980. The ecology and behavior of the Long-Billed Curlew in Southeastern Washington. Wildlife Monographs, No. 73.
- Apps, C.D. and N.J. Newhouse. 2000. Habitat modeling for badgers in the East Kootenay region of British Columbia. Columbia Basin Fish and Wildlife Compensation Program, Nelson, BC; East Kootenay Environmental Society, Kimberley, BC; Forest Renewal British Columbia, Cranbrook, BC; Parks Canada, Radium Hot Springs, BC.
- Apps, C.D., N.J. Newhouse and T.A. Kinley. 2001. Habitat associations of American Badgers in southeast British Columbia. Columbia Basin Fish & Wildlife Compensation Program.
- \_\_\_\_\_. 2002. Habitat associations of American badgers in southeastern British Columbia. Canadian Journal of Zoology 80:1228-1239.
- BC Environment. 1996. Managing Identified Wildlife Guidebook 1.0, Kamloops Forest Region. Ministry of Environment, Lands and Parks and Ministry of Forests. Internal Government Review Draft.
- \_\_\_\_\_. 1997. Species and Plant Community Accounts for Identified Wildlife. Ministry of Environment, Lands and Parks and Ministry of Forests. Internal Government Review Draft.
- \_\_\_\_\_. 1998. Habitat Atlas for Wildlife at Risk: South Okanagan and Lower Similkameen. Ministry of Water, Lands, and Air Protection, Penticton, BC.
- \_\_\_\_\_. 1999. Identified Wildlife Management Strategy: Volume 1. Produced by the Ministry of Environment, Lands, and Parks and Ministry of Forests, Victoria, BC.
- Beaucher, M.A. and J.A. Dulisse. 2004. First confirmed breeding record for the western screech-owl (*Megascops kennicottii macfarlanei*) in southeastern British Columbia. Northwest Naturalist 85:128-130.
- Bent, A.C. 1968. Life Histories of North American Cardinals, Grosbeaks, Buntings, Towhees, Finches, Sparrows, and Allies. Part 2. U.S. National Museum Bulletin 237: 1209-1217.
- Bertram, N., K.W. Larsen and J. Surgenor. 2001. Identification of critical habitats and conservation issues for the Western Rattlesnake and the Great Basin Gopher Snake within the Thompson-Nicola region of British Columbia. Prepared for the Ministry of Water, Land and Air Protection, Kamloops, BC and the Habitat Conservation Trust Fund of BC, Victoria, BC.
- Bishop, C.A., A.M. Bezener and R.J. Cannings. 2005. Draft National Recovery Strategy for the British Columbia Population of the Western Yellow-breasted Chat (*Icteria virens auricollis*). National Recovery Plan No. XX. Recovery of Nationally Endangered Wildlife. Ottawa. XX pp.
- Blood, D.A. and M. Macartney. 1998. Painted Turtle; Wildlife at Risk in British Columbia. Brochure. BC Environment. Victoria, BC.
- Bock, C.E., V.A. Saab, T.D. Rich and D.S. Dobkin. 1992. Effects of livestock grazing on neotropical migratory landbirds in western North America. In Status and Management of Neotropical Migratory Birds. USDA,

- Forest Service, Rocky Mountain Forest and Range Experiment Station, Colorado. General Technical Report RM-229.
- Brown, H.A., R.B. Bury, D.M. Darda, L.V. Diller, C.R. Peterson, and R.M. Storm. 1995. Reptiles of Washington and Oregon. Seattle Audubon Society, Seattle WA. 176 pages.
- Bryan, A. and L. Mulholland. 1992. Draft. Species Notes and Management Options for Fifty-four Wildlife Species of Management Concern in the South Okanagan. Ministry of Environment, Lands and Parks, Penticton, BC.
- Bryant, A.A. 1989. Late fall monitoring of Spotted Bats (*Euderma maculatum*) in the south Okanagan Valley, BC. Prepared for the Wildlife Branch, BC Environment. Penticton, BC.
- Campbell, R.W., A.K. Dawe, I. McTaggart-Cowan, J. Cooper, G. Kaiser and M.C. McNall. 1990. Birds of British Columbia: Volume 2, Nonpasserines; Diurnal Birds of Prey through Woodpeckers. Royal British Columbia Museum.
- Campbell, R.W., N.K. Dawe, I. MacTaggart-Cowan, J.M. Cooper, G.W. Kaiser, A.C. Stewart and M.C.E. McNall. 2001. The Birds of British Columbia, Volume 4: Passerines, Wood-warblers through Old World sparrows. Ministry of Environment, Lands and Parks, Victoria, BC, and Canadian Wildlife Service, Delta, BC.
- Cannings, R.A., R.J. Cannings and S.G. Cannings. 1987. Birds of the Okanagan Valley, British Columbia. Royal British Columbia Museum.
- Cannings, R.J. 1995. Status of the Grasshopper Sparrow in British Columbia. Wildlife Branch, Ministry of Environment Lands and Parks, Victoria, BC.
- \_\_\_\_\_. 1995. Status of the Yellow-breasted Chat in British Columbia. Wildlife Branch, Ministry of Environment Lands and Parks, Victoria, BC.
- \_\_\_\_\_. 1997. A survey of the Western Screech-Owl (*Otus kennicottii macfarlanei*) in the interior of BC. Prepared for Ministry of Environment, Lands and Parks.
- \_\_\_\_\_. 1998. COSEWIC status report on the Great Basin Spadefoot Toad (*Spea intermontana*) in Canada. Rep. prepared for the Committee on Endangered Wildlife in Canada. Ottawa, Ont.
- \_\_\_\_\_. 1999. Status of the Long-billed Curlew in BC. Ministry of Environment, Lands and Parks. Victoria, BC. Wildlife Working Paper No. WR-96.
- \_\_\_\_\_. 2004. Interior Western Screech-Owl, in Identified Wildlife management strategy – accounts and measures for managing Identified Wildlife, Version 2004. Ministry of Water, Land and Air Protection. Victoria, BC.
- Cannings, S.G., L.R. Ramsay, D.F. Fraser and M.A. Fraker. 1999. Rare Amphibians, Reptiles and Mammals of British Columbia. Ministry of Environment Lands, and Parks, Wildlife Branch and Resources Inventory Branch. Victoria, BC.
- CDC (Conservation Data Centre). 2005. Rare element occurrence database. Ministry of Sustainable Resource Management. Victoria, BC.
- Chapman, K., M.J. Sarell and the Southern Interior Amphibian Recovery Team. 2004. Draft National Recovery Plan for the Great Basin Spadefoot *Spea intermontana* in Canada. National Recovery Plan No. X. Recovery of Nationally Endangered Wildlife (RENEW). Ottawa, Ontario. XX pp.
- Charland, B. 1989. Size and winter survivorship in neonatal western rattlesnakes (*Crotalus viridis*). Canadian Journal of Zoology 67:1620-1625.
- Clarke, D., M. Martin, M. Sarell and C. Siddle. 1993. Greater Vernon Natural Areas/Features Inventory. Prepared for the Greater Vernon Parks and Recreation Department, Vernon, BC.

- Collard, T.S. and R.M.R. Barclay. 1991. Identification of the status and critical habitats of the Spotted Bat (*Euderma maculatum*), in the South Okanagan Valley, British Columbia. Unpublished report prepared for BC Environment. Victoria, BC.
- Collard, T.S., S.D. Grindal, R.M. Brigham and R.M.R. Barclay. 1990. Identification of the status and critical habitats of the Spotted Bat (*Euderma maculatum*), Pallid Bat (*Antrozous pallidus*) and Fringed Bat (*Myotis thysanodes*) in the South Okanagan Valley, BC. Unpublished report prepared for the Wildlife Branch, Ministry of Environment, Lands and Parks.
- Cooper, J.M. 1998. An inventory report on the status of diurnal raptors (Ferruginous Hawk, Swainson's Hawk, Prairie Falcon, Peregrine Falcon) at risk in the southern grasslands of British Columbia. Ministry of Environment, Lands and Parks. Victoria, BC. Wildlife Working Report WR-92.
- Deshant, J.A., L.M. Sondreal, D.H. Johnson, L.D. Igl, C.M. Goldade, P.A. Rabie, and B.R. Euliss. 2002. Effects of management practices on grassland birds: Long-billed Curlew. Northern Prairie Research Center, Jamestown, ND.
- De Smet, K.D. 1992. Status Report on the Long-billed Curlew (*Numenius americanus*) in Canada. Department of Natural Resources, Winnipeg, Man.
- Dobler, F.C. 1990. Shrub Steppe Ecosystem Studies. Phase One Completion Report. Nongame Research, Washington Department of Wildlife.
- Dorn, R.D. and J.L. Dorn. 1994. Further data on Screech-Owl distribution and habitat use in Wyoming. Western Birds. 25: 35-42.
- Easterla, D.A. 1973. Ecology of the 18 species of Chiroptera at Big Bend National Park, Texas. Northwest Missouri State University. Study 34: 1-165.
- Firman, M.C., C. Godwin and R.M.R. Barclay. 1995. Bat fauna survey of the West Shuswap and South Thompson River Region, British Columbia; Draft. Ministry of Environment, Lands and Parks. Victoria, BC.
- Fraser, D.F., W.L. Harper, S.G. Cannings and J.M. Cooper. 1999. Rare birds of British Columbia. Ministry of Environment, Lands and Parks, Wildlife Branch and Resources Inventory Branch. Victoria, BC.
- Fraser, D.F., T. Hooper and L.R. Ramsay. 1991. Preliminary Species Management Plan for Long-billed Curlew in British Columbia. Ministry of Environment Lands and Parks, Victoria, BC.
- Garcia, P.F.J., S.A. Rasheed and S.L. Holroyd. 1995. Status of the Spotted Bat in British Columbia. Ministry of Environment, Lands and Parks. Victoria, BC.
- Gebauer, M. 2004. Yellow-Breasted Chat, *in* Identified Wildlife management strategy – accounts and measures for managing Identified Wildlife, Version 2004. Ministry of Water, Land and Air Protection. Victoria, BC.
- Gibbard, B and M. Gibbard. 1992. Field Survey of the Yellow-breasted Chat in the South Okanagan. Ministry of Environment Lands and Parks, Penticton, BC.
- Green, D. M. and R. W. Campbell. 1984. The amphibians of British Columbia. Royal British Columbia Museum Handbook No. 45. Victoria, BC.
- Gregory, P.T. and R.W. Campbell. 1984. The Reptiles of British Columbia. Royal British Columbia Museum Handbook No. 44.
- Harvey, D.H. 1992. The Distribution, Density and Habitat of Brewer's Sparrows *Spizella breweri* in the South Okanagan Valley of British Columbia. Ministry of Environment Lands and Parks.
- Hayward, G.D. and O. Garton. 1984. Roost habitat selection by three small forest owls. Wilson Bulletin 96(4): 692-701.

- \_\_\_\_\_. 1988. Resource partitioning among forest owls in the River of No Return Wilderness, Idaho. *Oecologia* 75: 253-265.
- Hlady, D.A. 1990. South Okanagan Conservation Strategy. Ministry of Environment Lands and Parks, Victoria, BC.
- Hobbs, J. 2002. Confidential folio of Western Screech Owl Sites in the Southern Interior of BC. Habitat Branch, Ministry of Water, Land and Air Protection.
- Hobbs, J. and M. Sarell. 2001. An initial survey of the Identified Snakes in the Cariboo Forest Region. Prepared for FRBC and Ministry of Air, Water and Land Protection, Williams Lake, BC
- \_\_\_\_\_. Range of the Western Rattlesnake (*Crotalus viridis*) in British Columbia.
- Holroyd, S.L., R.M.R. Barclay, L.M. Merk and R.M. Brigham. 1994. A survey of the bat fauna of the dry interior of British Columbia. Ministry of Environment, Lands and Parks. Victoria, BC. Wildlife Working Report No. WR-63
- Holt, D.W. and J.M. Hillis. 1987. Current status and habitat associations of forest owls in Western Montana. Biology and conservation of Northern Forest Owls. Symposium Proceedings. US Department of Agriculture, Fort Collins, Colorado.
- Hoodicoff, C.S. 2003. Ecology of the Badger (*Taxidea taxus jeffersonii*) in the Thompson Region of British Columbia: implications for conservation. Master's thesis, University of Victoria. Victoria, BC.
- \_\_\_\_\_. 2005. Cariboo Region Badger project: 2004 field results and best management practices. Prepared for: Ministry of Water, Land and Air Protection, 100 Mile House, BC.
- Iverson, K. 2003. Sensitive Ecosystems Inventory: Bella Vista – Goose Lake Range. Volume 1: Methods, Ecological Descriptions, Results and Management Recommendations. Prepared for the Okanagan Indian Band, Allan Brooks Nature Centre, City of Vernon, and the Ministry of Water, Land and Air Protection.
- \_\_\_\_\_. 2005. Sensitive Ecosystems Inventory: Vernon Commonage. Volume 1: Methods, Ecological Descriptions, Results, Conservation Analysis and Management Recommendations.
- Iverson, K. and S. Shypitka. 2003. Sensitive Ecosystems Inventory: Bella Vista – Goose Lake Range. Volume 2: Terrestrial Ecosystem Mapping, Soil Erosion and Slope Stability, and Expanded Legends. Prepared for the Okanagan Indian Band, Allan Brooks Nature Centre, City of Vernon, and the Ministry of Water, Land and Air Protection.
- Iverson, K., and P. Uunila. 2005. Sensitive Ecosystems Inventory: Vernon Commonage. Volume 2: Terrestrial Ecosystem Mapping, Terrain Mapping, and Expanded Legend.
- jeffersonii* Badger Recovery Team. 2004. National Recovery Strategy for American Badger, *jeffersonii* subspecies (*Taxidea taxus jeffersonii*). Recovery of Nationally Endangered Wildlife (RENEW). Ottawa, Ontario.
- Kaufman, K. 1996. Lives of North American Birds (Peterson Natural History Companions). Houghton Mifflon Co., New York.
- Kinley, T.A. and N.J. Newhouse. 2005. East Kootenay Badger project 2004-2005 update: ecology, translocation, sightings and communications. Prepared for: Columbia Basin Fish and Wildlife Compensation Program, Nelson, BC, Parks Canada Agency, Radium Hot Springs, BC, and Ministry of Water, Land and Air Protection, Cranbrook, BC.
- Kirk, D.A. 1995. Status report on the Western Screech-Owl (*Otus kennicottii*) in Canada. Committee on the Status of Endangered Wildlife in Canada, Ottawa, Ont.
- Klauber, L.M. 1956. Rattlesnakes: their Habits, Life Histories, and Influence on Mankind. Vols. 1 & 2. University of California Press, Los Angeles, Ca.

- Knopp, D.H, A.E. Knopp and L.K. Larkin. 2000. Biophysical Survey of Vernon Military Camp. Prepared by BC's Wild Heritage Environmental Consultants, Sardis, BC. for the Department of National Defence, Canadian Forces Base Chilliwack.
- Krannitz, P.G. and C. Rohner. 2000. Habitat use and distribution of listed neotropical migrant songbirds in northeastern British Columbia. Pgs. 823-829 in L.M. Darling, ed. Proc. of a Conf. on the Biology and Management of Species at Risk. Wildl. Br., Ministry Environ., Lands and Parks and Univ. College of the Cariboo. February 1999, Kamloops, BC
- Krueper, D.J. 1992. Effects of land use practices on western riparian ecosystems. In Status and Management of Neotropical Migratory Birds. USDA, Forest Service, Rocky Mountain Forest and Range Experiment Station, Colorado. Gen.Tech.Rep. RM-229.
- Leonard, M.L. and M.B. Fenton. 1983. Habitat use by Spotted bats (*Euderma maculatum*, Chiroptera: Vespertilionidae): roosting and foraging behaviour. Canadian Journal of Zoology 61: 1487-1491.
- Leupin, E., D. J. Low and B. Persello. 1994. Census and life history observations of the Great Basin Spadefoot Toad (*Scaphiopus intermontanus*) breeding populations in the Thompson Nicola regions. Prepared for Wildlife Branch, BC Environment. Kamloops, BC.
- Lindzey, F.G. 1976. Characteristics of the natal den of the badger. Northwest Science 50: 178-180.
- Long, C.A. 1973. *Taxidea taxus*. Mammalian Species. No. 26:1-4.
- Low, B. 1976. The evolution of amphibian life histories in the desert. Pgs. 149-195 In D. W. Goodall (ed). Evolution of desert biota. University of Texas Press, Austin, Texas.
- Macartney M. 1985. The ecology of the Northern Pacific Rattlesnake, *Crotalus viridis oregonos*, in British Columbia. Master's thesis, University of Victoria, Victoria, BC. 289pp.
- Macartney, M., and P. T. Gregory. 1985. The Western Painted Turtle in Kikomun Creek Provincial Park. Unpublished report, submitted to Parks Branch, British Columbia.
- Marti, C.D. and J.S. Marks. 1987. Medium-sized owls. Proceedings of the Western Raptor Management Symposium and Workshop. National Wildlife Federation Scientific and Technical Series No. 12. Boise, Idaho.
- McAdoo, J.K., W.S. Longland and R.A. Evans. 1989. Nongame bird community responses to sagebrush invasion of crested wheatgrass seedings. Journal of Wildlife Management 53(2): 494-502.
- Messick, J.P. and M.G. Hornocker. 1981. Ecology of the Badger in southwestern Idaho. Wildlife Monographs 76.
- Ministry of Environment. 2005. Okanagan Region wildlife observation database. Fish and Wildlife Science and Allocation Section. Penticton, BC.
- Minta, S.C. 1993. Sexual differences in spatiotemporal interactions among badgers. Oecologia 96:402-409.
- Mylymok, J. and J. Hobbs. 2003. Inventory of Western Screech Owl (*O.k.macfarlanei*) in the Southern Interior of BC. Prepared for the BC Ministry of Water, Land and Air Protection, Penticton, BC.
- Nagorsen, D.W. and R.M. Brigham. 1993. Bats of British Columbia. Royal BC Museum Handbook. Victoria, BC.
- Nelson, K.J. and P.T. Gregory. 1992. A Survey of the Distribution, Biology and Population Trends of the Great Basin Gopher Snake, *Pituophis melanoleucus deserticola*, in British Columbia. Ministry of Environment Lands and Parks. Victoria, BC.
- Newhouse, N.J. and T.A. Kinley. 2000. Biology and conservation challenges of Badgers in the East Kootenay Region of British Columbia. In Proceedings of a Conference on the Biology and Management of Species and

- Habitats at Risk, Kamloops, BC, 15-19 Feb., 1999. Volume Two. L.M. Darling, editor. Ministry of Environment, Lands and Parks, Victoria, BC and University College of the Cariboo, Kamloops, BC.
- \_\_\_\_\_. 2000. Ecology of American Badgers near their range limit in Southeastern British Columbia. Columbia Basin Fish and Wildlife Compensation Program, Nelson, BC; Crestbrook Forest Industries, Cranbrook, BC; East Kootenay Environmental Society, Kimberley, BC; Parks Canada, Radium Hot Springs, BC.
- \_\_\_\_\_. 2004. East Kootenay Badger project 2003-2004 update: population ecology, translocation, sightings and communications. Prepared for: Columbia Basin Fish and Wildlife Compensation Program, Nelson, BC; and Parks Canada, Radium Hot Springs, BC.
- Nussbaum, R.A., E.D. Brodie, Jr. and R.M. Storm. 1983. Amphibians and Reptiles of the Pacific Northwest. University of Idaho Press, Moscow, Idaho.
- Ohanjanian, I. 1987. Status Report and Management Recommendations for the Long-billed Curlew (*Numenius americanus*) on the Junction. Ministry of Environment and Parks, Cariboo-Chilcotin Region, BC.
- Ohanjanian, I.A. 1992. Numbers, Distribution and Habitat Dynamics of Long-billed Curlews in the East Kootenay. Ministry of Environment, Cranbrook, BC.
- Ohanjanian, P. 2002. Long-billed Curlew in Standards for managing identified wildlife, Version 2. K. Paige ed. Ministry of Water, Land and Air Protection. Victoria, BC.
- Orchard, S.A. 1984. Amphibians and Reptiles of BC; an Ecological Review. Ministry of Forests. Victoria, BC.
- \_\_\_\_\_. 1985. Great Basin Spadefoot Toad (*Scaphiopus intermontanus*); Habitat Use Information and Habitat Suitability Index Model. Ministry of Forests, Victoria, BC.
- \_\_\_\_\_. 1988. Species notes on Reptiles, Vol. 3 of Wildlife habitat handbook for the Southern Interior Ecoprovince. BC Government, Victoria, BC.
- Orchard, S.A. and A.P. Harcombe. 1988. Species notes for reptiles. Volume 3 in A.P. Harcombe (tech. ed.). Wildlife habitat handbooks for the Southern Interior Ecoprovince. Ministry of Environment and Ministry of Forests. Victoria, BC. 47pp.
- Packham, R. 2004. Cariboo Region American Badger Project – Habitat Conservation Trust Fund project report. Ministry of Water, Land and Air Protection.
- Paczek, S. 2002. Grasshopper Sparrow (*Ammodramus savannarum perpallidus*). In Standards for managing identified wildlife, Version 2. K. Paige ed. Ministry of Water, Land and Air Protection. Victoria, BC.
- \_\_\_\_\_. 2002. Effects of fine scale and landscape level habitats features on sagebrush breeding birds of the south Okanagan and Similkameen valleys, BC. Unpublished Master's thesis, University of BC. Vancouver, BC.
- Pampush, G.J. 1980. Status Report on the Long-billed Curlew in the Columbia and Northern Great Basins. US Fish and Wildlife Service. Portland, Oregon.
- Pampush, G.J. and R.G. Anthony. 1993. Nest Success, Habitat Utilization and Nest-site Selection of Long-billed Curlews in the Columbia Basin, Oregon. The Condor. 95: 957-967.
- Rahme, A.H., A.S. Harestad and F.L. Bunnell. 1995. Status of the Badger in British Columbia. Ministry of Environment, Lands and Parks, Wildlife Branch. Victoria, BC. Wildlife Working Report WR 72.
- Resources Inventory Committee (RIC). 1998. Standard for Terrestrial Ecosystem Mapping in British Columbia. Ministry of Environment, Lands and Parks, Resource Inventory Branch. Victoria, B.C.  
<http://ilmbwww.gov.bc.ca/risc/pubs/tecolo/index.htm>

- \_\_\_\_\_. 1999. British Columbia wildlife habitat rating standards – Version 2.0. Ministry of Environment, Lands and Parks, Resource Inventory Branch. Victoria, BC. <http://ilmbwww.gov.bc.ca/risc/pubs/teecolo/index.htm>
- Royal British Columbia Museum. 1996. Endangered Species Web Site. Address: [http://rbcm1.rbcm.gov.bc.ca/End\\_Species/index\\_es.html](http://rbcm1.rbcm.gov.bc.ca/End_Species/index_es.html)
- Sarell, M. 1993. Snake hibernacula of the South Okanagan. Prepared for BC Environment, Penticton, BC and the BC Habitat Conservation Trust Fund, Victoria, BC.
- \_\_\_\_\_. 2004. Great Basin Spadefoot, *in* Identified Wildlife management strategy – accounts and measures for managing Identified Wildlife, Version 2004. Ministry of Water, Land and Air Protection. Victoria, BC.
- \_\_\_\_\_. 2004. Spotted Bat, *in* Identified Wildlife management strategy – accounts and measures for managing Identified Wildlife, Version 2004. Ministry of Water, Land and Air Protection. Victoria, BC.
- \_\_\_\_\_. 2005. Snake recovery strategy for the Vernon Military Cadet Camp. Prepared for the Chilliwack ASU, Department of National Defence, Chilliwack, BC.
- Sarell, M.J. and A. Haney. 2000. Rare Bats of the South Okanagan: 2000. Prepared for BC Environment, Penticton, BC.
- Sarell, M.J. and K.P. McGuinness. 1993. Rare bats of the shrub-steppe ecosystem of eastern Washington. Nongame Program. Department of Wildlife. Washington.
- Sarell, M.J., S. Robertson and A. Haney. 1997. Red and Blue-listed wildlife of the southern Boundary Forest District. FRBC Report.
- Sarell, M.J., S. Robertson and L. Scott. 1997. Inventory of snakes within Forest Development plans of the Boundary, Penticton, Merritt and Vernon Forest Districts. FRBC Report.
- Sargeant, A.B. and D.W. Warner. 1972. Movements and denning habits of a Badger. *Journal of Mammology* 53:207-210.
- Semlitsch, R.D. and J.R. Bodie. 2003. Biological criteria for buffer zones around wetlands and riparian habitats for amphibians and reptiles. *Conservation Biology* 17(5): 1219-1228.
- Schewchuk, C.H. 1996. The natural history of reproduction and movement patterns in the Gopher Snake (*Pituophis melanoleucas*) in Southern British Columbia. Master's thesis, University of Victoria. Victoria, BC.
- Shewchuk, C.H. and H.L. Waye. 1995. Draft. Status Report for the Gopher Snake in British Columbia. Ministry of Environment, Lands and Parks. Victoria, BC.
- Siddle, C. 1993. The Grasshopper Sparrow in the North Okanagan. Ministry of Environment Lands and Parks. Penticton, BC.
- \_\_\_\_\_. 1995. North Okanagan Long-billed Curlew Census. BC Environment. Penticton, BC.
- Smith, R.L. undated. *Ammodramus savannarum*: Grasshopper Sparrow. U.S. National Museum Bulletin 237. Plate 41.
- Stevens, V. 1995. Database for wildlife diversity in British Columbia: distribution and habitat use of amphibians, reptiles, birds and mammals in biogeoclimatic zones. Resource Branch, Ministry of Forests, Habitat Protection Branch, BC Environment, Victoria, BC. Working Paper 05/1995.
- St. Clair, R. C. 1989. The natural history of a northern turtle, *Chrysemys picta bellii* (Gray). Master's thesis, University of Victoria. Victoria, BC.

- St. John, D. 1993. Census of the breeding distribution of the Great Basin Spadefoot Toad, *Scaphiopus intermontanus*, in the south Okanagan Valley. Prepared for Wildlife Branch, BC Environment and Okanagan Region Wildlife Heritage Fund Society. Penticton, BC.
- Terres, J.K. 1995. The Audubon Society Encyclopedia of North American Birds. Wings Books, Avenel, New Jersey.
- Vander Haegen, W.M., F.C. Dobler and D.J. Pierce. 2000. Shrubsteppe bird response to habitat and landscape variables in Eastern Washington, USA. *Conservation Biology* 14(4): 1145-1160.
- Vickery, P.D. 1996. Grasshopper Sparrow (*Ammodramus savannarum*). In A. Poole and F. Gill, eds. The Birds of North America No. 239. Acad. Nat. Sci., Philadelphia, PA and Am. Ornithol. Union, Washington, DC.
- Wai-Ping, V. and M.B. Fenton. 1987. Habitat use and population status of *Euderma maculatum*, the Spotted Bat, in Southern British Columbia. Unpublished report prepared for the Nature Trust of BC.
- \_\_\_\_\_. 1989. Ecology of Spotted Bat (*Euderma maculatum*) roosting and foraging behaviour. *Journal of Mammology* 70(3): 617-622.
- Weins, J.A. 1973. Interterritorial Habitat Variation in Grasshopper and Savannah Sparrows. *Ecology* 54(4): 877-884.
- Weir, R.D., H. Davis and C. Hoodicoff. 2003. Conservation strategies for North American Badgers in the Thompson & Okanagan Regions. Final report for the Thompson-Okanagan Badger Project, Habitat Conservation Trust Fund.
- Weir, R.D., H. Davis, C.S. Hoodicoff, and K.W. Larsen. 2005. Life on a highway: Sources of mortality within an endangered British Columbian badger population. In T.D. Hooper, technical editor. Proceedings of the Species at Risk 2004 Pathways to Recovery Conference. March 2–6, 2004, Victoria, BC.
- Whitmore, R.C. 1981. Structural characteristics of grasshopper sparrow habitat. Short Communications. *Journal of Wildlife Management* 45(3).
- Yu, J.H.Y. 2001. Postfledging habitat use and movements of Brewer's Sparrows (*Spizella breweri breweri*) in the South Okanagan Region. Master's thesis, University of BC.

# Appendices

---

## Appendix A: Data Access

Spatial and non-spatial data for the Sensitive Ecosystems Inventory and Terrestrial Ecosystem Mapping (TEM), including wildlife mapping, are available for download at the former Ministry of Sustainable Resource Management's Terrestrial Ecosystem Mapping Data Warehouse at:

<http://srmwww.gov.bc.ca/ecology/tem/dataaware.html>

The following are available:

- Project metadata
- SEI report (Volume 1)<sup>32</sup>
- Arc/Info \*.E00 Export Files includes two spatial coverages: ECI field sampling points and a ECP TEM polygon coverage
- TEM Polygon Attributes
- TEM Map Legend Files
- TEM report with expanded legend (Volume 2)<sup>33</sup>
- Wildlife Species Accounts
- Wildlife Ratings Tables
- Wildlife Report (Volume 3)

---

<sup>32</sup> Iverson 2005

<sup>33</sup> Iverson and Uunila 2005

## Appendix B: Known and potential rare wildlife species in the study area.

Common Name	Scientific Name	Occurrence in Study Area	Prov. Status	COSEWIC Status
<b>Amphibians</b>				
Tiger Salamander	<i>Ambystoma tigrinum</i>	unknown	Red	Endangered
Great Basin Spadefoot	<i>Spea intermontana</i>	northern portion, likely throughout	Blue	Threatened
Western Toad	<i>Bufo boreus</i>	unknown but likely	-	Special Concern
<b>Reptiles</b>				
Painted Turtle	<i>Chrysemis picta</i>	throughout	Blue	-
Western Skink	<i>Eumeces skiltonianus</i>	unknown but possible	Blue	Special Concern
Western Rattlesnake	<i>Crotalus oreganus</i>	northern portion, likely throughout	Blue	Threatened
Gopher Snake	<i>Pituophis catenifer</i>	northern portion, likely throughout	Blue	Threatened
Racer	<i>Coluber constrictor</i>	northern portion, likely throughout	Blue	Special Concern
Rubber Boa	<i>Charina bottae</i>	throughout	-	Special Concern
<b>Birds</b>				
Great Blue Heron	<i>Ardea herodias</i> ssp. <i>herodias</i>	unknown but possible	Blue	-
California Gull	<i>Larus californicus</i>	known from one location	Blue	-
American Avocet	<i>Recurvirostre americana</i>	unknown but likely	Red	-
Long-billed Curlew	<i>Numenius americanus</i>	unknown but possible	Blue	Special Concern
Upland Sandpiper	<i>Bartramia longicauda</i>	unknown but possible	Red	-
Swainson's Hawk	<i>Buteo swainsoni</i>	northern portion, likely throughout	Red	-
Ferruginous Hawk	<i>Buteo regalis</i>	unknown but possible	Red	Special Concern
Western Screech-owl	<i>Megascops kennicotti</i> ssp. <i>macfarlanei</i>	historically (Ok Landing)	Red	Endangered
Flammulated Owl	<i>Otus flammeolus</i>	unknown but likely	Blue	Special Concern
Short-eared Owl	<i>Asio flammeus</i>	unknown but likely	Blue	Special Concern
White-throated Swift	<i>Aeronautes saxatalis</i>	northern portion	Blue	-
Lewis' Woodpecker	<i>Melanerpes lewis</i>	unknown but likely	Blue	Special Concern
Yellow-breasted Chat	<i>Icteria virens</i>	known from one location	Red	Endangered
Brewer's Sparrow	<i>Spizella breweri breweri</i>	unknown but possible	Red	-
Grasshopper Sparrow	<i>Ammodramus savannarum</i>	known from one location	Red	-
Lark Sparrow	<i>Chondestes grammacus</i>	known from one location	Red	-
<b>Mammals</b>				
Merriam's Shrew	<i>Sorex merriami</i>	unknown but possible	Red	-
Preble's Shrew	<i>Sorex prebeii</i>	unknown but possible	Red	-
Townsend's Big-eared Bat	<i>Corynorhinus townsendii</i>	unknown but likely (Bellavista)	Blue	-
Spotted Bat	<i>Euderma maculatum</i>	known from one location	Blue	Special Concern
Pallid Bat	<i>Antrozous pallidus</i>	unknown but possible	Red	Threatened
Fringed Myotis	<i>Myotis thysanodes</i>	unknown (Ok Landing)	Blue	Special Concern
Western Small-footed Myotis	<i>Myotis ciliolabrum</i>	unknown but likely	Blue	-
Western Harvet Mouse	<i>Reithrodontomys megalotis</i>	unknown but likely (Bellavista)	Blue	Special Concern
Great Basin Pocket Mouse	<i>Perognathus parvus</i>	unknown but likely	Blue	-
Nuttall's Cottontail	<i>Sylvilagus nuttallii</i> ssp. <i>nuttallii</i>	unknown	Blue	Special Concern
Badger	<i>Taxidea taxus</i>	northern portion, likely throughout	Red	Endangered



## Appendix D: Ratings Table

Ratings Table filename: Com wl ratings\_21Nov05.csv (See Appendix A for access)

Example of Ratings Table format:

ECO_SEC	BGC_ZONE	BGC_SUBZON	BGC_VRT	SITEMC_S	SITE_MA	SITE_MB	STRCT_S	STRCT_M	STAND_A	SERAL	A-SPIN_RE	A-SPIN_LIA	RCHPI_LIA	RCHPI_RE	R-CROR_LIS	R-CROR_LIA	R-PICA_LIG	R-PICA_RE	B-SWHA_RE	B-SWHA_LIG	B-LBCU_RE	B-LBCU_LIG	B-WSOW_RE	B-YBCH_LIG	B-GRSP_LIG	M-EUMA_RB	M-TATA_LIA
NOB	IDF	xh	1	AS			3				L	L	N	N	H	N	M	N	N	N	N	N	N	M	N	N	N
NOB	IDF	xh	1	AS			4		B		L	L	N	N	H	N	M	N	N	N	N	N	N	H	N	N	N
NOB	IDF	xh	1	AS			5		B		L	L	N	N	H	N	M	N	N	N	N	N	L	H	N	N	N
NOB	IDF	xh	1	AS			6		B		L	L	N	N	H	N	M	N	N	N	N	N	M	H	N	N	N
NOB	IDF	xh	1	AS			7		B		L	L	N	N	H	N	M	N	N	N	N	N	M	H	N	N	N
NOB	IDF	xh	1	AS	g		3				L	L	N	N	H	N	M	N	N	N	N	N	N	M	N	N	N
NOB	IDF	xh	1	AS	g		4		B		L	L	N	N	H	N	M	N	N	N	N	N	N	H	N	N	N
NOB	IDF	xh	1	AS	g		5		B		L	L	N	N	H	N	M	N	N	N	N	N	L	H	N	N	N
NOB	IDF	xh	1	AS	g		6		B		L	L	N	N	H	N	M	N	N	N	N	N	M	H	N	N	N
NOB	IDF	xh	1	AS	g		7		B		L	L	N	N	H	N	M	N	N	N	N	N	M	H	N	N	N
NOB	IDF	xh	1	AS	g	k	5		B		N	N	N	N	M	N	L	N	N	N	N	N	L	H	N	N	N
NOB	IDF	xh	1	AS	g	w	3				N	L	N	N	H	N	M	N	N	N	N	N	N	M	N	N	N
NOB	IDF	xh	1	AS	g	w	4		B		N	L	N	N	H	N	M	N	N	N	N	N	N	H	N	N	N
NOB	IDF	xh	1	AS	g	w	5		B		N	L	N	N	H	N	M	N	N	N	N	N	L	H	N	N	N
NOB	IDF	xh	1	AS	g	w	6		B		N	L	N	N	H	N	M	N	N	N	N	N	M	H	N	N	N
NOB	IDF	xh	1	AS	g	w	7		B		N	L	N	N	H	N	M	N	N	N	N	N	M	H	N	N	N
NOB	IDF	xh	1	AS	k		3				N	N	N	N	M	N	L	N	N	N	N	N	N	M	N	N	N
NOB	IDF	xh	1	AS	k		4		B		N	N	N	N	M	N	L	N	N	N	N	N	N	H	N	N	N
NOB	IDF	xh	1	AS	k		5		B		N	N	N	N	M	N	L	N	N	N	N	N	L	H	N	N	N
NOB	IDF	xh	1	AS	k		6		B		N	N	N	N	M	N	L	N	N	N	N	N	M	H	N	N	N
NOB	IDF	xh	1	AS	k		7		B		N	N	N	N	M	N	L	N	N	N	N	N	M	H	N	N	N
NOB	IDF	xh	1	AS	n		4		B		N	L	N	N	H	N	M	N	N	N	N	N	N	H	N	N	N
NOB	IDF	xh	1	AS	w		3				N	L	N	N	H	N	H	N	N	N	N	N	N	M	N	N	N
NOB	IDF	xh	1	AS	w		4		B		N	L	N	N	H	N	H	N	N	N	N	N	N	H	N	N	N
NOB	IDF	xh	1	AS	w		5		B		N	L	N	N	H	N	H	N	N	N	N	N	L	H	N	N	N
NOB	IDF	xh	1	AS	w		6		B		N	L	N	N	H	N	H	N	N	N	N	N	M	H	N	N	N